



UNIVERSITI PUTRA MALAYSIA

**ENZYME AIDED PEELING AND MEMBRANE REMOVAL OF LOCAL
MANDARINS (CITRUS SUHUIENSIS)**

LIU FANNY.

FSMB 2004 21

**ENZYME AIDED PEELING AND MEMBRANE REMOVAL OF LOCAL
MANDARINS (*Citrus suhuiensis*)**

By

LIU FANNY

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science

May 2004



Specially Dedicated
To
My Late Father Liu Shu Ping
&
CT

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Master of Science

**ENZYME AIDED PEELING OF LOCAL MANDARIN (*CITRUS
SUHUIENSIS*) AND LOCAL MANDARIN SEGMENTS**

By

LIU FANNY

May 2004

Chairman: Associate Professor Azizah binti Osman, Ph.D.

Faculty: Food Science and Biotechnology

A combination of pectinases and cellulases are able to selectively alter the albedo and segment membrane structure of citrus fruits and, hence, aid the removal of the peel, adhering albedo layer and also the segment membrane. This study was carried out to determine the optimum conditions needed to peel local mandarins using pectinases (Peelzym® IV, Novozyme, Switzerland) and cellulases (Celluclast® 1.5L, Novozyme, Switzerland). The experiment variables were enzyme concentration, vacuum pressure and vacuum infusion time. In the first part of the experiment, the local mandarins were first scored from stem end to the blossom end followed by immersion in 1000 ml of enzyme solution at a set vacuum pressure and ambient temperature (27 ± 1 °C). Only one parameter was varied in any one experiment. The latter part of the experiment was carried out using Response Surface Methodology (RSM) to determine the optimum combinations of enzyme concentration, vacuum pressure and vacuum infusion time to aid enzymatic segment membrane removal. Echip software was employed in the experimental design, calculate equations and statistical analysis. Peelzym® IV at 0.4 % v/w, 650 mm Hg vacuum and 16 minutes of vacuum time were found to be optimal for peel removal.

The enzyme-peeled fruits were judged by the panellists using three different sensory tests to ascertain its appeal to consumers. A significant ($P < 0.05$) difference between enzyme-peeled and hand-peeled segments was found, with the panelists preferring the enzyme-peeled segments. Celluclast® 1.5L at 4.52 % v/w, vacuum pressure at 370 mm Hg for 9 minutes was found to be optimal. After segment membrane removal, the membraneless local mandarin segments were then placed in different concentrations of sugar solutions to gauge consumer acceptance. Different sugar concentrations were used to emulate commercially available canned mandarin segments. Although varying concentrations of sugar solutions were used, the colour, odour, firmness, presence of adhering segment membrane and segment integrity were not affected as there was no significant ($P < 0.05$) difference among the samples. It was also observed that local mandarins stored in 15 °Brix was the preferred sugar concentration. As an overview, enzyme-peeled segments were found to be much more appealing as it had a much more intense orange colour, was firm with no loss of segment integrity, hence, was very well accepted by the panelists. Thus, enzyme aided peeling has a great potential as an alternative method to replace conventional methods of peeling.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGUPASAN KULIT AND SELAPUT MEMBRAN LIMAU MADU
(*CITRUS SUHUIENSIS*) DENGAN BANTUAN PENGGUNAAN ENZIM**

Oleh

LIU FANNY

Mei 2004

Pengerusi: Profesor Madya Azizah binti Osman, PhD

Fakulti: Sains Makanan dan Bioteknologi

Kombinasi pektinase dan selulase telah didapati berupaya mengubah struktur “albedo” dan selaput membran segmen buah sitrus secara selektif dan seterusnya membantu dalam pengupasan kulit, lapisan “albedo” yang terlekat serta selaput membran segmen. Kajian ini dilaksanakan dengan tujuan mengenalpasti keadaan optimum yang diperlukan untuk mengupas kulit limau madu dengan menggunakan pektinase (Peelzym® IV, Novozyme, Switzerland) dan selulase (Celluclast® 1.5L, Novozyme, Switzerland). Pembolehubah eksperimen adalah kepekatan enzim, tekanan vakum, serta tempoh infusi vakum. Pada mulanya, limau madu dikeluar kulitnya, dari hujung tangkai ke hujung pangkal, tanpa menempuhi bahagian isi limau. Langkah ini diikuti dengan perendaman limau dalam 1000 ml larutan enzim pada tekanan enzim yang ditetapkan pada suhu bilik (27 ± 1 °C). Hanya satu parameter diubahsuai dalam setiap satu langkah eksperimen. Bahagian kedua eksperimen dilaksanakan dengan menggunakan Methodologi “Response Surface” (RSM) untuk menentukan titik optimum kombinasi kepekatan enzim, tekanan vakum serta tempoh infusi vakum dalam pengupasan selaput membran segmen limau madu. Perisian lembut Echip digunakan untuk menentukan rekabentuk eksperimen, mengira keseimbangan dan membuat penilaian statistik. Keputusan menunjukkan bahawa

Peelzym® IV pada kepekatan 0.4 % v/w, 650 mm Hg vakum dan tempoh infusi enzim selama 16 minit adalah keadaan optimum bagi pengupasan kulit limau madu. Limau madu yang telah dibuang kulitnya dinilai oleh para panel dengan menggunakan tiga jenis kaedah sensori bagi menentukan penerimaan dan daya tarikannya oleh pengguna. Perbezaan yang ketara ($P < 0.05$) diantara limau yang dikupas tangan berbanding dengan limau yang telah dikupas dengan bantuan enzim telah didapati di mana para panel telah menunjukkan yang limau yang telah dikupas dengan bantuan enzim menjadi pilihan mereka. Didapati juga bahawa kepekatan enzim Celluclast® 1.5L pada kepekatan 4.52 % v/w, tekanan vakum pada 370 mm Hg serta tempoh infusi vakum selama 9 minit adalah keadaan optimum bagi pengupasan selaput segmen limau madu. Selepas pengupasan selaput segmen, segmen limau madu telah direndam di dalam larutan gula pada kepekatan yang berlainan untuk mengenalpasti penerimaan para pengguna. Langkah ini dilakukan untuk menyamai produk limau yang boleh didapati secara komersil. Walaupun kepekatan larutan gula yang berlainan telah digunakan, ciri fisiko-kimia segmen limau madu tidak terjejas. Didapati tiada perbezaan yang ketara ($P < 0.05$) ke atas warna, bau, kekerasan, kehadiran selaput yang masih terlekat serta integriti segmen juga masih kekal. Didapati juga produk hasil rendaman dalam larutan gula pada 15 °Brix merupakan pilihan para panel. Secara amnya, boleh dikatakan bahawa limau madu yang telah dibuang kulit dengan bantuan enzim adalah lebih menarik memandangkan ia mempunyai warna yang lebih menarik serta ketara, tidak terlalu lembut teksturnya serta diterima dengan baik oleh para pengguna. Dengan itu boleh disimpulkan bahawa teknik pengupasan kulit dan selaput segmen limau madu dengan bantuan enzim berpotensi untuk mengambil alih cara pengupasan konvensional yang lain.

ACKNOWLEDGEMENTS

I would like to convey my sincere appreciation and gratitude to the chairperson of my supervisory committee, Associate Professor Dr Azizah Osman for her all the advice, guidance, encouragement, personal concern and constant support. She has been an embodiment of kindness from the very beginning of my struggles and confusion to this time and space where I am about to graduate from this programme. Thank you for all that you have done and for your confidence in me.

I would also like to voice my sincere thanks to Professor Dr Hasanah M. Ghazali who was the one who brought about the decision for me to pursue my Master of Science in this field. I am truly grateful for the opportunity that has come about from this. Prof Hasanah was the one who brought about my awareness in the wonders of enzyme. She has been an inspiration ever since the day I attended her first lecture and reinforced since I started work on this project. Her encouragement throughout my research has really helped see me through a lot of hurdles and despairs. Her ability to constantly lead me to new wonders in research has really been a joy and an inspiration.

To Professor Dr Salmah Yusof, thank you for all your invaluable help in my research and the enjoyment of having worked with you especially in sensory evaluation. Your knowledge in this field and the advice you have provided is really appreciated.

To my group of true friends, Peng Kong, Yinsze, Kwan Kit and Chin Ping, I am really fortunate to have met all of you and bonded with you. Your encouragement and help throughout my research will never be forgotten. Special appreciation goes

to my lab mates, Dwi, Shita, Eja, and also Sook Meng, for making the lab a lively workplace . Thank you for all the laughter, the listening ear and the helpful hands.

Special appreciation to all the lab personnel, office staff and faculty members who has helped me in so many ways be it directly or indirectly. Thank you to each and everyone of you.

I would also like to express my gratitude and appreciation to my family members who has had the confidence in me, who has nurtured and educated me with the believe that I can achieve what I aim to be. Thank you for your unconditional love and support. A special appreciation goes to a very important person in my life, my father, who has always encouraged me to pursue my dreams. He has never failed to encourage me to go for my dreams, to reach out for as much as I can as the world is the limit when it comes to dreams. Although he is no longer here to see my achievements I know that he is and always will be happy and proud of all I have achieved and become.

Finally to a very important person in my life, Chi Tsung who has given me constant encouragement in completing this thesis, who has selflessly burned the midnight oil with me, words are insufficient to help me express my gratitude and appreciation. Thank you for your love and your faith in me. You never cease to amaze me with your actions and words. You have made my life very meaningful.

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

Azizah Osman, Ph.D.


Associate Professor,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia
(Chairman)

Hasanah Mohd Ghazali, Ph.D.

Professor,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia
(Member)

Salmah Yusof, Ph.D.

Professor,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia
(Member)



AINI IDERIS, Ph. D
Professor/Dean,
School of Graduate Studies,
Universiti Putra Malaysia.

Date: 10 SEP 2004

I certify that an Examination Committee met on 13th May 2004 to conduct the final examination of Liu Fanny on her Master of Science thesis entitled “Enzyme-aided Peeling and Membrane Removal of Local Mandarins (*Citrus suhuiensis*)” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Nazamid Saari, Ph.D.

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Chairman)

Azizah Osman, Ph.D.


Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

Hasanah Mohd. Ghazali, Ph.D.

Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

Salmah Yusof, Ph.D.

Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **26 AUG 2004**

DECLARATION

I hereby declare that the thesis is based on my original work except for the quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or any other institutions.


LIU FANNY

Date: 18 Aug 2004

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF PLATES	xviii
 CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	6
2.1 Citrus fruits	
2.1.1 Citrus	7
2.1.2 Citrus fruits in Malaysia	13
2.1.3 Mandarins	17
2.1.4 Local mandarins	20
2.2 Physico-chemical characteristics of local mandarins	24
2.2.1 Fruit Composition	25
2.2.2 Total Soluble Solids (TSS)	26
2.2.3 Sugar Content	28
2.2.4 Organic Acids and pH	28
2.2.5 Ascorbic Acid (Vitamin C)	30
2.2.6 Peel Colour	30
2.2.7 Pectins	32
2.2.8 Pectic Substances	33
2.3 Fruit peeling	34
2.3.1 Conventional Methods of Fruit Peeling	34
2.3.2 Pectic Enzymes	38
2.3.3 The Use of Enzyme Infusion For Peeling	45
2.3.4 Citrus Peeling	48
2.3.5 Future Uses of Enzyme Infusion	49
3 PHYSICO-CHEMICAL CHARACTERISTICS OF LOCAL MANDARINS	51
3.1 Introduction	51
3.2 Materials and methods	52
3.2.1 Source of fruits	52
3.2.2 Sampling	52
3.2.3 Physical Analyses	52
3.2.4 Chemical Analyses	54

3.2.5	Statistical Analysis	56
3.3	Results and discussion	56
3.3.1	Physical Analyses	56
3.3.2	Chemical Analyses	59
3.4	Conclusion	61
4	EFFECTS OF ENZYME CONCENTRATION VACUUM INFUSION TIME AND VACUUM PRESSURE ON ENZYME AIDED PEEL REMOVAL	63
4.1	Introduction	63
4.2	Materials and methods	65
4.2.1	Local mandarins	65
4.2.2	Sample Preparation	66
4.2.3	Enzyme Preparation	66
4.2.4	Effect of Enzyme Concentration	66
4.2.5	Effect of Vacuum Intensity	67
4.2.6	Effect of Vacuum Time	67
4.2.7	Peeling Quality and Efficiency Criteria	67
4.2.8	Sensory Evaluation	68
4.3	Results and discussion	70
4.3.1	Effect of Enzyme Concentration	70
4.3.2	Effect of Vacuum Intensity	73
4.3.3	Effect of Vacuum Time	76
4.3.4	Sensory Evaluation	80
4.4	Conclusion	89
5	EFFECTS OF ENZYME CONCENTRATION VACUUM PRESSURE AND VACUUM INFUSION TIME ON MEMBRANE REMOVAL OF LOCAL MANDARINS	91
5.1	Introduction	91
5.2	Materials and methods	93
5.2.1	Source of local mandarins	93
5.2.2	Sample Preparation	93
5.2.3	Enzyme Preparation	94
5.2.4	Experimental Design	94
5.2.5	Quality and Efficiency Criteria for Membrane Removal	95
5.2.6	Sensory Evaluation	97
5.3	Results and discussion	100
5.4	Conclusion	116
6	CONCLUSION AND RECOMMENDATIONS	117
	BIBLIOGRAPHY	119
	APPENDICES	127
	BIODATA OF THE AUTHOR	147

LIST OF TABLES

Table	Page
2.1.1 Chemical characteristics of local mandarin juice	23
2.1.1a Composition of California Valencia orange juice and local mandarin juice	24
2.2.1 Nutrient composition of different citrus fruits	25
2.2.1a Nutrient composition of local mandarin	27
2.2.2 The available carbohydrates of fruits	29
2.2.3 Approximate levels of vitamin C in some fruits and vegetables	31
2.3.1a Cellulases, hemicellulases and pectinases in food biotechnology	40
2.3.1b Cellulases, hemicellulases and pectinases in food biotechnology	41
3.1 Pulp-to-peel ratio, number of segments and percent edible portion of three different batches of local mandarins	57
3.2 Mean values of total soluble solids (TSS), pH, ascorbic acid (Vitamin C), titratable acidity, Hunter L, 'a and 'b, Fructose, glucose and sucrose, pectin in pulp and skin contents of three batches (replicates) of local mandarins	58
4.1 Effect of different enzyme concentrations for peel removal of local mandarins (<i>Citrus suhuiensis</i>)	71
4.2 Effect of different vacuum pressures needed for peel removal of local mandarins (<i>Citrus suhuiensis</i>)	74
4.3 Effects of different vacuum time needed for peel removal of local mandarins (<i>Citrus suhuiensis</i>)	78
4.4 Mean of preference scores (hedonic scores) for colour, odour, firmness, sliminess, appearance and overall acceptability between hand peeled and enzyme peeled local mandarin segments	88
5.1 Combinations of enzyme concentration, vacuum pressure and vacuum time needed for complete membrane removal of local mandarins	96
5.2 Response of overall incubation time based on the trials generated by Echip on the variables of enzyme concentration, vacuum time and vacuum pressure needed for segment membrane removal of local mandarins	101

5.3	Regression Coefficients, R^2, and P or Prob. Values^a for the overall incubation time needed for enzymatic removal of local mandarin segment membrane	102
5.4	Average Hedonic scores for various attributes of commercial mandarin segments	107
5.5	Average Hedonic scores for sensory evaluation of enzyme-peeled local mandarin segments treated with 5 different sugar concentrations	114

LIST OF FIGURES

Figure	Page
2.1.1 Interspecific and intergeneric hybrids of citrus	8
2.1.2 Cross section of a citrus food	10
2.1.3 Common products of citrus processing industry	14
2.1.4 Products from Malaysian citrus	17
2.2.1 Structure of pectin molecule	34
2.3.1 Enzymatic mode of action of the most frequent depolymerases on the pectin molecule	43
4.1 Percent ability to detect between sample different/ odd sample and same sample	81
4.2 Percentage of difference between hand-peeled and enzyme-peeled segments	83
4.3 Percent preference between enzyme-peeled and hand-peeled segments	83
4.4 Average intensity scores of panelists for colour, odour, firmness, sliminess, appearance and overall acceptability between hand peeled and enzyme peeled local mandarin segments	86
4.5 Percent preference between enzyme-peeled and hand-peeled segments of local mandarins	88
5.1 Contour map on the effects of enzyme concentration, vacuum time and vacuum pressure on the overall incubation time of segment membrane removal	104
5.2 QDA evaluation of the attributes of three different commercial samples of peeled, canned mandarins for the selection of standard	106
5.3 Percent preference between enzyme-peeled lab samples and commercial mandarin samples	110
5.4 Survey data on the degree of difference between enzyme-peeled lab samples and commercial samples conducted at the end of the triangle test	110
5.5 Score distribution of enzyme-peeled local mandarin segments in 5 different sugar concentrations	112

5.6	Percent preference for enzyme-peeled local mandarins treated with 5 different sugar concentrations	115
-----	--	-----



LIST OF PLATES

Plate		Page
1	Local mandarin (<i>Citrus suhuiensis</i>)	21
2	Local mandarin (<i>Citrus suhuiensis</i>) tree	22
3	Enzyme-peeled whole fruit (<i>Citrus suhuiensis</i>)	82
4	Hand-peeled whole fruit (<i>Citrus suhuiensis</i>)	82
5	Enzyme-peeled local mandarin segments	84
6	Hand-peeled local mandarin segments	84
7	Enzyme-peeled membraneless segments	108
8	Hand-peeled membraneless segments	108

CHAPTER 1

INTRODUCTION

Fruit and fruit products, in all its many varieties and variations, are major world commodities and part of the economic life-blood of many countries, particularly the developing world (Arthey and Ashurst, 1996). Fruits have long been valued as part of the staple diet of many living things. As a matter of fact, fruits have been inexorably linked to the welfare of life on our planet (Taylor, 1996). Fruits are a form or class of food that is not only affordable but delicious, easily obtainable and highly nutritious. Most fruits, are primarily water, but also contain over 400 other constituents and nutrients, i.e. carbohydrates, organic acids, amino acids, ascorbic acid, minerals, flavonoids, carotenoids, volatiles, lipids, etc.

Nutrients frequently consumed in sub-optimal concentrations by humans are proteins, iron, vitamin A, thiamin (Vitamin B₁), riboflavin (Vitamin B₂) and ascorbic acid (Fourie 1996). Members of this group are often known as the critical nutrients. Fruits also contain these critical nutrients, and sometimes some of these nutrients occur in higher concentrations than in other foods (Fourie, 1996). Frequent consumption of vegetables and fruits, especially green and yellow vegetables and citrus fruits, is associated with decreased susceptibility to some forms of cancer, although the mechanisms for their protective effects have not been fully determined. Therefore, fruits play an important role in balancing the human diet, mainly because the compositions of fruits differ markedly from other foods of plant and animal

origins (Hugo, 1969). Fruits are often consumed fresh or processed, as a result of the many attributes of the many different varieties of fruits available. Therefore, it is inevitable that processing of fruits are important to maximize the utility of fruit and to encourage an increase in fruit consumption.

Peeling is one of the most important preparatory steps in the processing of some fruits and vegetables meant for canning, freezing and dehydration (Radhakrishnaiah-Setty *et al.*, 1993) and also as a fresh refrigerated item. In the beginning of the fruit processing industry, only hand peeling was practiced. Over the years, several other methods of peeling i.e. peeling by steam, mechanical peeling, cryogenic peeling, etc were introduced, improving the efficiency of peeling. With the advancement in technology, enzyme (pectinases and cellulases) -aided processing (peeling, segmentation and membrane removal) has been introduced. This new method of peeling has many advantages over the other conventional methods/ practices of peeling. Among the advantages are minimization of production losses, energy and chemical use, heat ring formation and pollution load, and in addition it peels to the extent dictated by the products (Radhakrishnaiah-Setty *et al.*, 1993).

The United States Department of Agriculture (USDA) patented (USDA Patent No. 4,284,651) enzyme infusion technology in 1981 (Bruemmer, 1981) but it is relatively new to Malaysia. It has not been thoroughly studied especially in the context of local citrus. Local fruits i.e. citrus have been found to be very high in nutrients and

fiber. Moreover this is a good way of promoting local fruits, presented in a different form to local consumers.

Enzyme treatment offers a way to achieve very specific changes in whole foods or food ingredients. Moreover, the removal or inactivation of the enzyme requires no specific label reference to enzyme use making it very consumer friendly (McArdle and Culver, 1994). Interest in enzyme infusion has increased in proportion to the availability and purity of commercial enzymes (McArdle and Culver, 1994). Consumers are beginning to show an interest in less-processed, fresher tasting retail food products (especially by the catering and bakery industry) that are fresh, convenient, can be prepared and consumed in less time.

The availability and purity of commercial enzymes have also contributed to the interest in enzyme infusion. In addition to that, the load of pollution caused by lye peeling can be greatly reduced when using enzyme infusion. This will be of particular importance, as consumers are becoming more and more conscious of the importance of non-environmental polluting products.

In addition, “natural” alteration of food ingredients has been a major rationale for genetic manipulation of plants (Caransa *et al.*, 1998). Consumer concerns over genetically engineered food ingredients (Fraley, 1992) may produce a climate where alternative biological strategies such as enzyme infusion accomplish the same task as

genetic engineering in a simpler scheme. Enzyme infusion may also lead to unique opportunities, such as improved processes for peeling and sectioning, changes that may be difficult or impossible through genetic manipulation (McArdle and Culver, 1994).

In recent years, consumers have become more health-conscious in their food choices but have less time to prepare healthful meals. As a result, the market demand for 'minimally processed' fruits and vegetables has rapidly increased (Jacxsens *et al.*, 1999). Enzyme infusion can be driven not only by the food industry's interest in responding to consumer needs but also the need to increase productivity.

Malaysia is rich in resources and has an abundant supply of local fruits that are mostly non-seasonal (easily available), cheap and highly nutritious. These resources are under-utilised. Therefore, it is important to realise the potential of using high technological methods of processing these fruits minimally. With this, Malaysia might be able to penetrate into the international food market as a major producer of processed and minimally processed fruits. There is great potential in texture-improved ready-to-eat citrus fruits with intact fruit sacs and also for separated fruit sacs that can be incorporated into fruit juice in Malaysia.

The local mandarin / "Limau Madu" (*Citrus suhuiensis*) is a local citrus that is easily obtainable and available. With the aid of enzymes, it is relatively easy to process

citrus fruits to produce attractive, value added, peeled fruits. These minimally processed citrus would be in demand in the catering industry such as the airline, hotel and bakery industries as it is less time consuming and more efficient as compared to hand peeling (the conventional method used in Malaysia).

Therefore the objectives of this study were to:

1. determine and compare the physico-chemical characteristics of local mandarins with other citrus fruits.
2. determine the effects of enzyme concentration, vacuum pressure and vacuum time needed for optimum peel removal of local mandarins.
3. optimize enzyme aided membrane removal of local mandarins with RSM (Response Surface Methodology).
4. determine consumer acceptance of enzymatically peeled (removal of skin and segment membrane) local mandarins.

All the data obtained will be useful to determine the optimum conditions needed for peel and membrane removal in order to optimize processing and to reduce production cost. This is in the belief that this new technology has great potential as an alternative to conventional peeling.

CHAPTER 2

LITERATURE REVIEW

2.1 Citrus fruits

2.1.1 Citrus

Citrus is considered as one of the important fruit crops in tropical and sub-tropical regions. According to Davies and Albrigo (1994), the general area of origin of citrus is believed to be southeastern Asia, including that from eastern Arabia east to the Philippines and from the Himalayas south to Indonesia or Australia. Within this large region, northeastern India and northern Myanmar were believed to be the center of origin, but recent evidence suggests that Yunnan Province in south-central China may be as important due to the diversity of species found and the system of rivers that could have provided dispersal to the south (Gmitter and Hu, 1990). Extensive movement of the various types of citrus probably occurred within the general area of citrus origin from before recorded history. Currently, citrus is grown primarily between the latitudes 40°N to 40°S. More northern and southern locations of commercial production exist where temperatures are moderated by ocean winds (Davies and Albrigo, 1994).

Production of fresh and processed citrus surpasses that of bananas, apples, grapes, or mangoes (Davenport, 1990). Fresh citrus has been appreciated to the extent that it was carried from its area of origin and cultivated from before the fifth century BC